

Genotypic variation in leaf epicuticular wax quantity in a large faba bean (*Vicia faba* L.) germplasm collection

Hamid Khazaei^{1,2*}, Arja Santanen¹, Kenneth Street³, and Frederick L. Stoddard¹

¹ Department of Agricultural Sciences, Viikki Plant Science Centre, Helsinki Sustainability Science Centre, University of Helsinki, Helsinki, Finland, ² Current address: Department of Plant Sciences, University of Saskatchewan, Saskatoon, Canada and ³ International Center for Agricultural Research in the Dry Areas (ICARDA), Rabat, Morocco

*Corresponding author: H. Khazaei, ORCID ID: 0000-0002-5202-8764, contact:

hamid.khazaei@gmail.com, call @ +1 (306) 361-1167

Running title: **Epicuticular wax diversity in faba bean**

Published in Plant Genetic Resources: Characterization and Utilization (2019) 17: 298-300. Doi: 10.1017/S1479262118000461

Abstract

Among grain legumes, faba bean is reputed to be relatively sensitive to drought stress. Epicuticular wax (ECW) quantity is considered as an important drought adaptation strategy in plant species. This study aimed to define variation in leaf ECW concentration as a drought-adaptive trait in 197 faba bean accessions under well watered conditions. The relationship between ECW and stomatal characteristics was also investigated. Highly significant differences were found in the ECW concentration, which ranged from 0.680 to 2.104 mg/dm². No relationships were found between ECW and any measure of stomatal morphology and function. This study provides evidence of wide variation in ECW in faba bean germplasm, which is independent of stomatal characteristics and leaf water content. This variation may allow the genetic improvement of ECW as a drought-adaptive character in faba bean breeding programs aiming at economical use of water.

Keywords: epicuticular wax, genetic diversity, faba bean, drought, stomata

Introduction

Faba bean is an excellent source of protein for human food and animal feed, and it is suitable for environmentally sustainable agriculture due to its generous ecological services. Nevertheless, faba bean is reputed to be relatively sensitive to drought (McDonald and Paulsen, 1997; Daryanto *et al.*, 2015). Various morpho-physiological and biochemical characters confer drought adaptation in plants. Several studies have demonstrated faba bean drought-related morpho-physiological traits in above-ground (*e.g.*, Darwish and Fahmy, 1997; Khan *et al.*, 2010) and below-ground characteristics (*e.g.*, Belachew *et al.*, 2018). The waxes on plant cuticles form a protective cover against biotic and abiotic stresses (Jenks and Ashworth, 1999; Xue *et al.*, 2017). The effectiveness of stomatal control over transpiration efficiency increases if non-stomatal transpiration pathways are restricted. An important non-stomatal water vapour pathway is the leaf cuticle.

Accumulation of epicuticular wax (ECW) is considered an important drought avoidance strategy in many plant species. In pea (*Pisum sativum* L.), a greater amount of epicuticular wax was correlated with a higher harvest index under drought conditions (Sánchez *et al.*, 2001). Drought stress enhanced the deposition of cuticular waxes and subsequently improved drought adaptation in alfalfa (*Medicago sativa* L., Zhang *et al.*, 2005) and soybean (*Glycine max* (L.) Merr., Kim *et al.*, 2007). Variation in the wax quantity has been demonstrated in several crop species (reviewed in Sharma *et al.*, 2018) but not yet in faba bean. Thus, this study aimed to explore variation in leaf ECW concentration as a drought adaptation character in a relatively large faba bean germplasm collection.

Experimental

A set of 197 faba bean accessions from the ICARDA collection was used in this study (Table S1).

Faba bean germplasm was grown in 2 L plastic pots containing a mixture of sand and peat (3:1 v/v) in a climate-controlled glasshouse at the Department of Agricultural Sciences, University of Helsinki, as reported previously (Khazaei *et al.*, 2013). The experimental design was a randomized complete block with four replicates. Throughout the experiment, soil moisture level was maintained at field capacity. Photoperiod was adjusted to 14 h/10 h (light/dark), and the temperature was maintained at 21°C day/15°C night. Minimum daytime PPFD was set to 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at the canopy level. Relative humidity was kept at 60%.

Eight weeks after sowing, the following measurements were taken on the youngest, fully expanded leaves. Leaflet area was measured using a LI-6200 leaf area meter (LI-COR Inc., Lincoln, NE). Stomatal density and area were measured following the impression method (Wang and Clarke 1993). Gas exchange traits were measured using a LI-6400 portable photosynthesis system. Canopy temperature was measured using a FLUKE® 574 infrared gun (Fluke Corp., Everett, WA). Leaf relative water content was measured as described by Barrs and Weatherley (1962).

The amount of ECW was determined by a colorimetric wax quantification method developed by Ebercon *et al.* (1977), based on the colour change produced by the reaction of ECW with $\text{K}_2\text{Cr}_2\text{O}_7$. The reagent was prepared by mixing 20 g of powdered potassium bichromate with 40 ml of deionized water. The slurry was mixed with 1 L of concentrated sulfuric acid and heated at 80°C until a clear solution was obtained. To extract wax, 10 fresh leaf discs having a total area of 40 cm² (both surfaces) were punched from the leaves. Samples were

74 immersed in 17 ml of chloroform for 15 s. The chloroform-ECW solution was filtered through a
75 fine filter paper, then evaporated on a boiling water bath until the odour of chloroform was gone.
76 Next, 5 ml of $K_2Cr_2O_7$ reagent was added into glass test tubes and heated at 100°C in a water
77 bath for 30 min. After cooling, 12 ml of deionized water was added, and the colour was allowed
78 to develop for at least 3 h. The optical densities of the samples (two readings per replicate) were
79 measured at 590 nm with a PharmaSpec UV-1700 UV–Visible Spectrophotometer (PharmaSpec,
80 Shimadzu, Germany).

81 For reference purposes, faba bean wax was extracted from a large amount of field-grown
82 leaves using chloroform. A standard curve was prepared by following a serial dilution method
83 from ECW extracted from PEG-3000 (polyethylene glycol), Carnauba wax, and faba bean wax.
84 The linear standard curve equation was used to determine the wax concentration of samples.

85 The R statistical package ([R Development Core Team, 2016](#)) was used for all data
86 analysis. The means of studied traits are given in [Table S2 and S3](#).

Results and Discussion

The standard curves for all three tested waxes were linear throughout the concentrations (Fig. S1). PEG-3000 was curved slightly downward and Carnauba slightly upward while faba bean wax was straight. The faba bean wax was used as the standard for ECW measurements.

The 197 accessions differed for ECW ($P < 0.001$). ECW ranged from 0.680 mg/dm² to 2.104 mg/dm² (Fig. 1; Table S2), with a mean \pm SD of 1.288 ± 0.263 mg/dm². ECW showed no significant correlations with stomatal morphology, gas exchange traits and relative water content, and only a weak positive association with leaflet area ($P < 0.090$) (Table 1). The lack of associations between ECW and other leaf traits related to drought adaptation, particularly stomatal characteristics under non-stress conditions, indicates that faba bean might use both rapid stomatal closure and increased wax accumulation to reduce water loss when adapting to drought. In stress conditions, however, ECW was associated with the gas exchange traits in *Jatropha* (Figueiredo *et al.*, 2012) and wheat (Huggins *et al.*, 2018). It should be noted that leaf mesophyll conductance, chloroplast density, and cell wall thickness also regulate water balance in plants (Ouyang *et al.*, 2017).

Here we employed a high-throughput screening colorimetric method for screening a large number of faba bean accessions for ECW. To our knowledge, no previous study has shown the variability in ECW in faba bean. Accessions IG 12132, IG 12260, IG 11989, IG 137686, IG 72375, IG 112096 and IG 74554 had the highest ECW concentrations among accessions. These accessions might be used in genetic improvement of ECW as a component of drought adaptation in faba bean breeding programs.

108 **Acknowledgments**

109 H.K. expresses his gratitude to CIMO (Centre for International Mobility) and the Emil Aaltonen
110 Foundation for their financial support during the experiments.

References

- Barrs HD and Weatherley PE (1962) A re-examination of the relative turgidity technique for estimating water deficit in leaves. *Australian Journal of Biological Sciences* 15: 413–428.
- Belachew KY, Nagel KA, Fiorani F, Stoddard FL (2018) Diversity in root growth responses to moisture deficit in young faba bean (*Vicia faba* L.) plants. *PeerJ* 6: e4401.
- Darwish DS and Fahmy GM (1997) Transpiration decline curves and stomatal characteristics of faba bean genotypes. *Biologia Plantarum* 39: 243–249.
- Daryanto S, Wang L and Jacinthe P-A (2015) Global synthesis of drought effects on food legume production. *PLoS One* 10: e0127401.
- Ebercon A, Blum A and Jordan WR (1977) A rapid colorimetric method for epicuticular wax content of sorghum leaves. *Crop Science* 17: 179–180.
- Figueiredo KV, Oliveira MT, Oliveira AFM, Silva GC and Santos MG (2012) Epicuticular-wax removal influences gas exchange and water relations in the leaves of an exotic and native species from a Brazilian semiarid region under induced drought stress. *Australian Journal of Botany* 60: 685–692.
- Huggins T, Mohammed S, Sengodan P, Ibrahim AMH, Tilley M and Hays DB (2018) Changes in leaf epicuticular wax load and its effect on leaf temperature and physiological traits in wheat cultivars (*Triticum aestivum* L.) exposed to high temperature during anthesis. *Journal of Agronomy and Crop Science* 204 :49–61.
- Jenks MA and Ashworth EN (1999) Plant epicuticular waxes: function, production and genetics. In: Janick, J. (ed.) *Horticultural Reviews*. John Wiley & Sons, Inc. Oxford, UK. pp. 1–68.
- Khan HR, Paull JG, Siddique KHM and Stoddard FL (2010) Faba bean breeding for drought affected environments: A physiological and agronomic perspective. *Field Crops Research* 115: 279–286.

134 Khazaei H, Street K, Bari A, Santanen A and Stoddard FL (2013) Do faba bean (*Vicia faba* L.) accessions
135 from environments with contrasting seasonal moisture availabilities differ in stomatal
136 characteristics and related traits? *Genetic Resources and Crop Evolution* 60: 2343–2357.

137 Kim KS, Park SH, Kim DK and Jenks MA (2007) Influence of water deficit on leaf cuticular waxes of
138 soybean (*Glycine max* [L.] Merr.). *International Journal of Plant Sciences* 168: 307–316.

139 McDonald GK and Paulsen GM (1997) High temperature effects on photosynthesis and water relations of
140 grain legumes. *Plant and Soil* 196: 47–58.

141 Ouyang W, Struik PC, Yin X and Yang J (2017) Stomatal conductance, mesophyll conductance, and
142 transpiration efficiency in relation to leaf anatomy in rice and wheat genotypes under drought.
143 *Journal of Experimental Botany* 68: 5191–5205.

144 R Development Core Team (2016) R: A language and environment for statistical computing. R
145 foundation for statistical computing, Vienna, Austria. ISBN 3–900051–07–0. Available at:
146 <http://www.R-project.org>.

147 Sánchez FJ, Manzanares M, de Andrés EF, Tenorio JL and Ayerbe L (2001) Residual transpiration rate,
148 epicuticular wax load and leaf colour of pea plants in drought conditions. Influence on harvest
149 index and canopy temperature. *European Journal of Agronomy* 15: 57–70.


150 Sharma P, Kothari SL, Rathore MS and Gour VS (2018) Properties, variations, roles, and potential
151 applications of epicuticular wax: a review. *Turkish Journal of Botany* 42: 135–149.

152 Wang H and Clarke JM (1993) Genotypic, intraplant, and environmental variation in stomatal frequency
153 and size in wheat. *Canadian Journal of Plant Science* 73: 671–678.

154 Xue D, Zhang X, Lu X, Chen G and Chen Z-H (2017) Molecular and evolutionary mechanisms of
155 cuticular wax for plant drought tolerance. *Frontiers in Plant Science* 8: 621.

156 Zhang JY, Broeckling CD, Blancaflor EB, Sledge MK, Sumner LW and Wang ZY (2005)
157 Overexpression of *WXPI*, a putative *Medicago truncatula* AP2 domain-containing transcription
158 factor gene, increases cuticular wax accumulation and enhances drought tolerance in transgenic
159 alfalfa (*Medicago sativa*). *The Plant Journal* 42: 689–707.

160 **Figure legends**



161
 162 **Fig. 1.** Frequency distribution of leaf epicuticular wax concentration in 197 faba bean accessions
 163 (Skewness= 0.316, Kurtosis= 0.474). $LSD_{0.05}$ (least significant differences) = 0.059.

164 **Table 1.** Pearson correlation coefficients of epicuticular wax content (ECW) with morpho-
 165 physiological traits among 197 faba bean accessions.

	LA	SD	SA^a	g_s	E	TE^b	CT	RWC
Correlation with ECW	0.121	0.099	-0.088	0.030	0.042	0.000	-0.040	0.029

166 ^a Stomatal length × stomatal width.

167 ^b Transpiration efficiency was calculated as photosynthetic rate divided by the stomatal conductance.

168 *LA*, leaflet area; *SD*, stomatal density; *SA*, stomatal area; *g_s*, stomatal conductance; *E*, transpiration rate;

169 *TE*, transpiration efficiency; *CT*, canopy temperature; *RWC*, relative water content.